

Warm-up Problems - April , 2006

Solutions

1. Find the probabilities for the standard normal distribution.

(a) $P(0 \leq X \leq 2.26) = 0.4881$

(b) $P(X \leq 2.26) = 0.9881$

(c) $P(X \geq 2.26) = 0.0119$

(d) $P(-2.26 \leq X \leq 2.26) = 0.4881 + 0.4881 = 0.9762$

(e) $P(1.08 \leq X \leq 2.26) = 0.3599 + 0.4881 = 0.8480$

(f) $P(X \leq -2.26) = 0.0119$

Lecture Problems

2. Convert the X -probabilities into Z -probabilities.

Find the probabilities for the normal distribution indicated.

(a) $\mu = 2.5, \sigma = 3$

$$P(X \geq 4) = P(Z \geq 0.5) = 0.3085$$

(b) $\mu = -4, \sigma = 2.1$

$$P(-3.5 \leq X \leq -3) = P(0.24 \leq Z \leq 0.48) = 0.1844 - 0.0948$$

(c) $\mu = 6.2, \sigma = .12$

$$P(X \leq 5.95) = P(Z \leq -2.08) = 0.0188$$

(d) $\mu = 6.2, \sigma = 2.1$

$$P(X \leq 5.95) = P(Z \leq -0.12) = 0.4522$$

3. Course scores are normally distributed with $\mu = 65$ and $\sigma = 5$.

- (a) Find the proportion of students who scores above 70.

Solution:

$$P(X \geq 70) = P(Z \geq 1) = 0.1587$$

- (b) Find the proportion of students who scored above 80.

Solution:

$$P(X \geq 80) = P(Z \geq 3) = 0.0013$$

- (c) If the top 10% of the class receive A's, find the lowest score to receive an A.

Solution: Here we find the z score with $P(Z \geq z) = 0.1$, which gives us $z = 1.28$. We now find the corresponding x -score:

$$x = \mu + z\sigma = 65 + (1.28)(5) = 71.40$$

- (d) If the top 40% of the class receive A's and B's, find the lowest score to receive a B.

Solution: Here we find the z score with $P(Z \geq z) = 0.4$, which gives us $z = 0.25$. We now find the corresponding x -score:

$$x = \mu + z\sigma = 65 + (0.25)(5) = 66.25$$

Area Under the Standard Normal Curve ($\mu = 0, \sigma = 1$)
From 0 to z

[illegible]