

Math 233 - November 13, 2009

- More versions of Green's Theorem

$$\begin{aligned}\oint_{\partial R} p \, dx + q \, dy &= \iint_R \frac{\partial q}{\partial x} - \frac{\partial p}{\partial y} \, dA \\ &= \iint_R \text{Rot} \, (p, q) \, dA \\ &= \int_{\partial R} F \cdot \mathbf{u} \, ds \\ \oint_{\partial R} F \cdot \mathbf{n} \, ds &= \iint_R \text{Div} \, F \, dA\end{aligned}$$

1. Let $r(t) = (t, t^3)$.
 - (a) Find a tangent vector to the curve for any time t .
 - (b) Find a vector normal to the curve for any time t .
 - (c) Does your normal vector point to the right or to the left of the direction of the curve. Find the normal vector in the other direction too.
 - (d) Find both unit normal vectors
2. A wire in the shape of a semi-circle, $r(t) = (3 \cos t, 3 \sin t)$, $0 \leq t \leq \pi$. has density equal to $\delta(x, y) = y$ (in other words, the wire is heavier as y increases). Using an integral $\int_C \delta(x, y) \, ds$, find the mass of the wire.

Lecture Problems

3. Let $F = (x^2 + y^2, xy)$ and C be the boundary of the square with vertices $(0, 0)$, $(1, 0)$, $(1, 1)$ and $(0, 1)$. Using Green's Theorem, calculate the circulation of F around C .
4. Let $F = (-y, x)$ and let r be the circle of radius 2 center at the origin. Find the flux of F across r .
5. Let $F = (2y, xy)$. Find the flux across the unit circle.
6. Use Green's Theorem to compute the flux of the vector field $F = (2x, xy)$ across the unit circle.