

Math 233 - October 26, 2009

- What is so great about $G = \left(-\frac{y}{x^2+y^2}, \frac{x}{x^2+y^2}\right)$?
 - How can we extend the idea of integration to $\mathbb{R}^2$?
1. True/False. If false, determine how the question can be made true.
 - (a) The work done by a vector field around a loop is zero.
 - (b) If $P, Q \in \mathbb{R}^n$ then the work that vector field F does from P to Q is the negative of the work done from Q to P .
 - (c) Suppose F is not conservative and $r(t)$ a path from P to Q . Then the work that F does along r from P to Q is equal to the negative of the work done along the reverse path from Q to P .
 - (d) Suppose the curl of the vector field F is equal to $(0, 0, 0)$. Then the word done around every loop is equal to 0.
 2. Find the line integrals using potential functions.
 - (a) Let $r(t)$ be the line from $(1, 5)$ to $(-2, 3)$.
$$\int_r y^2 dx + (2xy - 1) dy =$$
 - (b) Let $r(t) = (1 + \cos t, -2 + 3 \sin t)$ for $0 \leq t \leq \pi/2$.
$$\int_r (2xy^3 + y - 1) dx + (3x^2y^2 + x) dy =$$
 - (c) Let $r(t)$ be the upper hemisphere of the unit circle oriented counter-clockwise.
$$\int_r (y - e^y) dx + (x - xe^y) dy =$$

Lecture Problems

3. Let $r(t)$ be the unit circle oriented positively. Compute

$$\frac{1}{2\pi} \oint \frac{y^3 + x^2y - 4x}{2(y^2 + x^2)^2} dx - \frac{(xy^2 + 4y + x^3)}{2(y^2 + x^2)^2} dy =$$

4. Let

$$G = \left(-\frac{y}{x^2 + y^2}, \frac{x}{x^2 + y^2}\right)$$
$$F = \left(\frac{y^3 + x^2y - 4x}{2(y^2 + x^2)^2}, -\frac{(xy^2 + 4y + x^3)}{2(y^2 + x^2)^2} dy\right)$$

Find k and ϕ so that $F = kG + \nabla\phi$.

$$k =$$
$$\nabla\phi =$$
$$\phi =$$